



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

Update prepared by
Climate Prediction Center / NCEP
June 10, 2013



Outline

- Overview
- Recent Evolution and Current Conditions
- MJO Index Information
- MJO Index Forecasts
- MJO Composites



Overview

- The MJO has remained fairly weak over the past several days, with the enhanced phase centered over the eastern Indian Ocean.
- Dynamical model MJO index forecasts are in reasonable agreement for a strengthening MJO signal during the next week, with eastward propagation to the Western Pacific by the end of two weeks.
- Based on recent observations and model MJO forecasts, the MJO is forecast to strengthen and become slightly better organized across the Maritime Continent over the next two weeks.
- Enhanced rainfall is favored during Week-1 across parts of southern Asia, and the Maritime Continent. Tropical cyclogenesis is more likely than normal over the South China Sea, north of the Philippines, and over the western Caribbean Sea.
- During Week-2 the MJO favors enhanced convection across parts of the southeast Asia and the Maritime Continent.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies have become more organized over Africa while easterly anomalies decreased over the equatorial Indian Ocean.

Easterly anomalies persisted over most of the western Pacific.

Westerly anomalies continued over the East Pacific during the past five days.



850-hPa Zonal Wind Anomalies (m s^{-1})



Westerly anomalies (orange/red shading)
represent anomalous west-to-east flow

Easterly anomalies (blue shading)
represent anomalous east-to-west flow

Westward propagation (dashed/solid lines sloping down and to the left) of anomalies during early December were primarily due to equatorial Rossby wave activity as the MJO was then generally weak.


During late December the MJO strengthened (alternating dotted/dashed lines).

During March and early April, anomalies indicate signs of being influenced by equatorial Rossby wave activity with less eastward propagation evident.

The MJO strengthened during early May with eastward propagation of low-level wind anomalies noted. More recently, other sub-seasonal modes have limited eastward propagation.




OLR Anomalies – Past 30 days




Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During early May, convection rapidly developed across the Indian Ocean, while the central Pacific experienced suppressed convection.



During mid-May, the OLR field became less coherent with generally enhanced convection north of the Equator across the Indian Ocean and portions of the Maritime Continent. Convection also increased in parts of the East Pacific due to a northward displaced ITCZ




Most recently in late May, the OLR field has remained incoherent with respect to the MJO. A convective dipole across the Indian Ocean and anomalous convection across Central America appear to be the dominant features.



Outgoing Longwave Radiation (OLR)

Anomalies (7.5°S-7.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

The MJO was a dominant mode of variability across the Tropics from January into March as indicated by the alternating dashed and dotted lines.

Near the end of March, the anomalies show signs of influence from other modes of tropical variability. However, MJO activity reemerged in early April across the Indian Ocean.


During early May, OLR decreased significantly (stronger negative anomalies) across the Indian Ocean. The MJO signal broke down during the middle of the month and remains fairly weak, with other modes of variability influencing the pattern.



200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation
Negative anomalies (green shading) indicate favorable conditions for precipitation

Time ↓



The MJO strengthened in late December, (alternating dashed and dotted lines) and anomalies increased in magnitude with more robust eastward propagation indicated during late 2012 to April 2013.

Anomalies became less coherent at times during late January and early February as the influence from other modes of variability are evident in the depicted anomalies. Some reorganization is evident in late February and early March.


The velocity potential anomalies were more coherent only briefly during early and mid-May and have since broken down, with other modes of tropical variability influencing the pattern.



IR Temperatures (K) / 200-hPa

Velocity Potential Anomalies


Positive anomalies (brown contours) indicate unfavorable conditions for precipitation
Negative anomalies (green contours) indicate favorable conditions for precipitation



The velocity potential pattern suggests some modest reorganization of the MJO with the convectively active phase across the Indian Ocean. However, other modes remain important as the MJO signal is weak.



200-hPa Vector Wind Anomalies (m s^{-1})



Note that shading denotes the zonal wind anomaly


Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Large-scale tropical anomalies remain fairly weak, with westerly anomalies over much of the central Pacific.



200-hPa Zonal Wind Anomalies (m s^{-1})



Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow


Eastward propagation of westerly wind anomalies associated with the MJO is evident beginning in late December and continuing into April 2013. Some propagation of easterly anomalies is evident during late January and early February.

During March and early April, anomalies were influenced by westward moving features over the central and western Pacific.

Westerly anomalies shifted east of the Date Line during early May. The rapid phase speed suggests the influence of a higher-frequency Kelvin waves. Recently, westerly anomalies have become persistent over the central and eastern equatorial Pacific.



Weekly Heat Content Evolution in the Equatorial Pacific



Through August 2012, heat content anomalies became positive and increased in magnitude across the eastern equatorial Pacific, partly in association with a downwelling Kelvin wave.

An oceanic Kelvin wave was initiated at the end of September and increased heat content across the central and eastern Pacific during October and November.

Positive (negative) anomalies developed in the western (eastern) Pacific during January 2013 and persisted into early March. The influence of a downwelling oceanic Kelvin wave can be seen during late February and March as anomalies became positive in the east-central Pacific.

Recently, the pattern is persistent, with some cooling east of 115W.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index indicates some strengthening of the MJO signal in Phase 4 over the past several days. However, the MJO remains weak.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.
Plots put current MJO activity in historical context.




Ensemble GFS (GEFS) MJO Forecast

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days
light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The bias-corrected ensemble GFS indicates a strengthening MJO signal, with some propagation through Phase 4. A weakening of the signal is implied during Week-2.

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The ensemble mean GFS forecasts weakly enhanced convection across the Indian Ocean and Maritime Continent with some suppressed convection across the east Pacific during week-1. Later, some eastward and northward propagation of the enhanced convection is forecast over south Asia.




Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




This statistical forecast indicates some strengthening over the Indian Ocean during Week-1, with some propagation eastward and northward during Week-2.




MJO Composites – Global Tropics

850-hPa Velocity Potential and Wind Anomalies (May-Sep)



Precipitation Anomalies (May-Sep)






U.S. MJO Composites – Temperature

- Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9


<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>