



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
November 18, 2013**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- The MJO continued to become less coherent during the previous week, with influence from other types of coherent tropical subseasonal variability dominating the anomalous convection and circulation pattern.
- Dynamical model MJO index forecasts indicate little signal over the next several days as other types of subseasonal variability continue to strongly influence the tropical circulation. The majority of dynamical MJO index forecasts do show the potential for a weak eastward moving signal re-emerging over the eastern Indian Ocean and Maritime Continent during Week-2. It is too early to conclude that this potential signal will emerge as a longer lived, more robust MJO.
- Statistical forecasts suggest a continued weak MJO signal.
- Based primarily on the latest observations and some dynamical model guidance, the MJO is not forecast to contribute significantly to anomalous tropical convection at the current time. Any influence from an emerging MJO would contribute to enhanced convection over the Maritime Continent and northern Australia.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Easterly anomalies persisted over the western Indian Ocean, while a small area of westerly anomalies developed over the eastern Indian Ocean and western Maritime Continent.

Easterly anomalies continued across the western Pacific while spreading east of the Date Line.

Westerly anomalies persisted over the eastern Pacific during the past five days, and increased over western Africa.




850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time

↓



Longitude

The MJO strengthened during June and continued until mid-July with fast eastward propagation.

During late July through mid-August, the MJO was weak. In late August and early September, westerly (easterly) anomalies increased over the eastern (western) Pacific in associated with renewed MJO activity.

During October, equatorial Rossby wave activity was strong from 160E to 100E as westward movement features are evident (red box). MJO activity was less coherent during this period.


Westerly anomalies persisted across the Western Hemisphere during late October and early November, with strong influence from westward moving features evident over the Maritime Continent and Indian Ocean.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)



During late October, enhanced convective activity was observed over the western and eastern Pacific north of the equator, associated with tropical cyclone activity. Enhanced (suppressed) convection was observed over South Asia (Philippines and South China Sea).


During early November, enhanced convection associated with tropical cyclones continued over the western Pacific. Enhanced (suppressed) convection was observed over South America (eastern Indian Ocean)

During mid-November, suppressed (enhanced) convection was observed over the western Pacific (Maritime Continent). Enhanced convection developed over the Horn of Africa and the equatorial Indian Ocean, while suppressed convective anomalies expanded across parts of South America.



Outgoing Longwave Radiation (OLR)

Anomalies (7.5°N-7.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

The MJO strengthened once again during June and continued into early July.

MJO was active during late August and September with the enhanced phase propagating eastward over the western Pacific Ocean, while the suppressed phase strengthened over the Indian Ocean.

Tropical cyclone activity contributed to the persistence of enhanced convection across the West Pacific as well as a weakened suppressed phase further west.

Recently there has been some development of convective anomalies across the western Indian Ocean with a fast eastward propagation.

Longitude




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time



Longitude

The MJO was active (alternating dashed and dotted lines) during June and early July before weakening at the end of the month.

The MJO was not active during late July and much of August, but strengthened during late August and September, with eastward propagation of robust upper-level velocity potential anomalies. Other modes of tropical intraseasonal variability are also evident.


During the second half of October, upper-level velocity potential exhibited little MJO related variability. Recently, some coherent eastward propagation on the fast side of the MJO envelope of phase speeds has been observed, although the signal started to break down during early November.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The velocity potential pattern has become less coherent, with generally anomalous large scale upper level divergence over Africa, the Indian Ocean, and Maritime Continent, and anomalous large scale upper-level convergence over the Atlantic, South America, and the western Pacific. Influence from a possible Kelvin wave is evident over the central Pacific.




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly upper-level zonal wind anomalies increased over the equatorial western Pacific and eastern Pacific. Easterly anomalies diminished over the tropical Atlantic.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow




The MJO strengthened (alternating dotted and dashed lines) during June and its influence continued to mid-July, as eastward propagation of wind anomalies associated with the MJO were again observed.

During August, westerly wind anomalies were generally persistent just west of the Date Line. Renewed MJO activity occurred during late August and September with westerly wind anomalies shifting east to the eastern Pacific.

Most recently, anomalies of alternating sign have continued over the eastern Pacific, due in part to extratropical Rossby waves breaking into the Tropics.



Weekly Heat Content Evolution in the Equatorial Pacific



An oceanic downwelling Kelvin wave was initiated at the end of September and increased heat content across the central and eastern Pacific during October and November 2012.

Positive (negative) anomalies developed in the western (eastern) Pacific during January 2013 and persisted into early March. The influence of a downwelling oceanic Kelvin wave can be seen during late February and March as anomalies became positive in the east-central Pacific.

Evidence of oceanic downwelling Kelvin waves are seen in late August and October/November.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.


Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).





MJO Index -- Recent Evolution

- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months





MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.
Plots put current MJO activity in historical context.



Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members


Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The ensemble GFS indicates an incoherent MJO signal under the influence of other modes of variability during Week-1, with eastward propagation of a weak signal over the Maritime Continent during Week-2.







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The ensemble mean GFS forecasts little anomalous convection during Week-1, with enhanced convection over the eastern Indian ocean propagating over the Maritime Continent during Week-2.



Constructed Analog (CA) MJO Forecast


Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



The constructed analog MJO forecast indicates a weak signal.


Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days






MJO Composites – Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov-Mar)



Precipitation Anomalies (Nov-Mar)






U.S. MJO Composites – Temperature

- Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>