



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
December 30, 2013**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- Although the MJO remained weak during the past week, some diagnostic tools indicate a coherent pattern with the enhanced convective phase over the Maritime Continent. Other types of coherent subseasonal tropical variability also remained active.
- A large spread exists among the dynamical model MJO index forecasts regarding the evolution of the MJO during the next two weeks.
- The MJO is forecast to contribute to enhanced (suppressed) convection over parts of the western South Pacific (Indian Ocean) during the next two weeks. Enhanced probabilities for tropical cyclone development are forecast for the Indian Ocean during Week-1 and across the South Pacific during the next two weeks.
- Enhanced convection along and south of the equator in the West Pacific due in part to the MJO could contribute to long wave amplification across the Pacific/North America region during Week-2 and beyond, enhancing the likelihood of cold air outbreaks across the central U.S.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly anomalies diminished over the equatorial Indian Ocean during the past five days while shifting eastward to the Maritime Continent.


Easterly anomalies persisted over the equatorial western Pacific.



850-hPa Zonal Wind Anomalies ($m s^{-1}$)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow
Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time
↓



Longitude

The MJO strengthened during June and continued until mid-July with fast eastward propagation.

During late July through mid-August, the MJO was weak. In late August and early September, westerly (easterly) anomalies increased over the eastern (western) Pacific in associated with renewed MJO activity.


During October, equatorial Rossby wave activity was strong from 160E to 100E as westward movement features are evident (red box). MJO activity was less coherent during this period.

During November and December, easterly anomalies have been persistent from 120E to near the Date Line. The easterly anomalies are smaller during the latter half of December.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)




Wet conditions were observed across the eastern Indian Ocean, southern Maritime continent, and northern Australia during late November and early December, while suppressed convection continued for much of the Pacific basin.

During early to mid-December, enhanced convection persisted across the Indian Ocean, while suppressed convection continued over much of the equatorial Pacific.

During mid to late December, enhanced convection shifted east to the Maritime Continent with suppressed convection persisting at the Date Line. Enhanced convection developed across the South Pacific.



Outgoing Longwave Radiation (OLR) Anomalies (17.5°S-2.5°S)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

The MJO was active from late August through early October (apparent in the Northern Hemisphere) with the enhanced phase propagating eastward from the Indian Ocean to the western Pacific Ocean over this period.

The MJO was generally weak or incoherent for much of November and other types of coherent tropical subseasonal variability were very active.

During late November, a large area of enhanced convection developed in the Indian Ocean and recently propagated eastward to the west Pacific Ocean.

Longitude




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time
↓



The MJO was active (alternating dashed and dotted lines) during June and early July before weakening at the end of the month.

The MJO was not active during late July and much of August, but strengthened during late August and September, with eastward propagation of robust upper-level velocity potential anomalies. Other modes of tropical intraseasonal variability are also evident.

From late October to early December, the MJO was not very strong or coherent. There was evidence of coherent eastward propagation at times during this period, but much of this activity exhibited fast propagation speeds more consistent with atmospheric Kelvin waves.

A slower eastward propagation of 200-hPa velocity potential anomalies was observed during mid-December.


Longitude



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The current velocity potential anomalies indicate a coherent pattern of large scale upper-level divergence (convergence) over the Maritime Continent and west Pacific (South America, Atlantic Ocean, and Africa).




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies spread eastward from the Pacific Ocean to northern South America during the previous five days (red box).



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time



Longitude


The MJO strengthened (alternating dotted and dashed lines) during June and its influence continued to mid-July, as eastward propagation of wind anomalies associated with the MJO were again observed.

During August, westerly wind anomalies were generally persistent just west of the Date Line. Renewed MJO activity occurred during late August and September with westerly wind anomalies shifting east to the eastern Pacific.

Anomalies of alternating sign are evident over the eastern Pacific, due in part to extratropical Rossby waves breaking into the Tropics (red bracket). Also, westerly anomalies have increased in December just east of the Date Line, while easterly anomalies developed over the Indian Ocean.



Weekly Heat Content Evolution in the Equatorial Pacific



Positive (negative) anomalies developed in the western (eastern) Pacific during January 2013 and persisted into early March. The influence of a downwelling oceanic Kelvin wave (dashed line) can be seen during late February and March as anomalies became positive in the east-central Pacific.

Oceanic downwelling Kelvin wave activity is evident in late August and once again during October through early December, the latter being the strongest wave to date in 2013.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution





- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index indicated an eastward propagation of a weak signal during the past week.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 1997 to present.
Plots put current MJO activity in historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The ensemble GFS forecast indicates a weak signal during Week-1 with an increase in amplitude during Week-2.







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days



The ensemble mean GFS forecasts enhanced convection persisting across the west/central Pacific during the next two weeks, while suppressed convection intensifies across the Indian Ocean later in Week-2.





Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days

Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days





The constructed analog MJO forecast indicates an eastward propagation of enhanced (suppressed) convection to the Western Hemisphere (Maritime Continent) during the next two weeks.



MJO Composites – Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov-Mar)

Precipitation Anomalies (Nov-Mar)






U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>