



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
February 10, 2014**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- The MJO remained incoherent during the past week, with influence from more localized subseasonal features such as tropical cyclone activity over the southwestern Indian Ocean, enhanced northern Australian monsoon activity, and suppressed convection over South America impacting the pattern of global tropical convection.
- There is considerable spread among the model guidance resolving the future evolution of the MJO, with several dynamical models indicating a continued weak signal and some statistical tools suggesting renewed MJO activity.
- Based on the latest observations and most model forecasts, the MJO is forecast to remain weak during the next 1-2 weeks, while lower frequency signals continuing to dominate the pattern.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly anomalies developed across southern Africa and parts the southwest Indian Ocean during the past five days.

Easterly anomalies expanded into the eastern Maritime Continent, while westerly anomalies persisted across the western Maritime Continent and northern Australia.

Westerly anomalies were observed across much of the equatorial Pacific east of the Date Line.




850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time

↓



Longitude

In late August and early September, westerly (easterly) anomalies increased over the eastern (western) Pacific in associated with renewed MJO activity.

During October, equatorial Rossby wave activity was strong from 160E to 100E as westward movement features are evident (red box). MJO activity was less coherent during this period.

During November and December, easterly anomalies were persistent from 120E to near the Date Line. Westerly anomalies were also evident across the Indian Ocean during this period.


During January, westerly anomalies intensified and shifted east to the Maritime Continent and West Pacific, but easterly anomalies over the Maritime Continent disrupted the signal during early February.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)



During mid-January, enhanced (suppressed) convection was observed across the Maritime Continent, West Pacific, and north central Australia (south central Indian Ocean, central Pacific, and eastern Australia).


Enhanced (suppressed) convection persisted across much of the western Pacific (eastern Indian Ocean) during late January. Enhanced convection was also observed across southern Africa and the southwest Indian Ocean, while suppressed convection persisted across southeastern Brazil.

During early February, enhanced convection persisted across parts of the southwest Indian Ocean and northern Australia, while suppressed convection developed across the Maritime Continent and persisted across eastern Brazil.



Outgoing Longwave Radiation (OLR)

Anomalies (7.5°N-7.5°S)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

The MJO was active from late August through early October with the enhanced phase propagating eastward from the Indian Ocean to the western Pacific Ocean over this period.

The MJO was generally weak or incoherent for much of November and other types of coherent tropical subseasonal variability were very active.

A large area of enhanced convection developed over the Indian Ocean during late November and propagated slowly eastward to the west Pacific Ocean by late January. This feature weakened during early February as suppressed convection propagated from the Indian Ocean to the Maritime Continent.




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time



Longitude

The MJO was not active during late July and much of August, but strengthened during late August and September, with eastward propagation of robust upper-level velocity potential anomalies (alternating dashed and dotted lines). Other modes of tropical intraseasonal variability are also evident.

From late October to early December, the MJO was not very strong or coherent. There was evidence of coherent eastward propagation at times during this period, but much of this activity exhibited fast propagation speeds more consistent with atmospheric Kelvin waves.


A slower eastward propagation of 200-hPa velocity potential anomalies was observed from mid-December to mid-January across the Indo-Pacific warm pool region (red box), while positive anomalies propagated from the Indian Ocean to the Maritime Continent during late January and early February.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The current velocity potential data indicates increasing organization of anomalies, with areas of large scale anomalous ascent (descent) over much of the Indian Ocean and northern Australia (Western Hemisphere). The velocity potential pattern remains largely disorganized over the Pacific.




200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time



Longitude

Renewed MJO activity (alternating dotted and dashed lines) occurred during late August and September with westerly wind anomalies shifting east to the eastern Pacific.

Anomalies of alternating sign are evident over the eastern Pacific, due in part to extratropical Rossby waves breaking into the Tropics (red bracket).

Westerly anomalies increased in December across the western Hemisphere and persisted into early January. Recently, anomalies have been dominated by Kelvin wave activity and interaction with the extratropics over the Western Hemisphere.




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly anomalies persisted across the east Pacific during the past ten days (red boxes).

Weak easterly (westerly) anomalies developed over the western (eastern) Indian Ocean during early February.



Weekly Heat Content Evolution in the Equatorial Pacific



Positive (negative) anomalies developed in the western (eastern) Pacific during January 2013 and persisted into early March. The influence of a downwelling oceanic Kelvin wave (dashed line) can be seen during late February and March as anomalies became positive in the east-central Pacific.

Oceanic downwelling Kelvin wave activity is evident in late August and once again during October through early December, the latter being the strongest wave during 2013.

There is evidence that another strong downwelling event is beginning in the West Pacific.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution




- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index remained weak during the previous 7 days, with some signal over the Maritime Continent.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 2007 to present.

Plot puts current MJO activity in recent historical context.



Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members


Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The ensemble GFS forecast indicates a weak MJO signal, with some redevelopment of convection over the Western Pacific by Week-2.







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The ensemble mean GFS forecasts convective anomalies over the Maritime Continent to propagate over the Western Pacific, with little additional eastward propagation during Week-2.



Constructed Analog (CA) MJO Forecast


Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



The constructed analog MJO forecast indicates a greater eastward propagation of the convective anomalies over the Pacific and Western Hemisphere during Week-2.


Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days






MJO Composites – Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov-Mar)



Precipitation Anomalies (Nov-Mar)






U.S. MJO Composites – Temperature

- Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>