



# **Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions**

**Update prepared by  
Climate Prediction Center / NCEP  
July 7, 2014**



# Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



# Overview

- **The MJO weakened during the past week as other modes of tropical subseasonal variability strongly influenced the pattern.**
- **Several dynamical model solutions indicate a strengthening MJO signal during the upcoming week, with the enhanced phase propagating over the Maritime Continent to the far western Pacific.**
- **Dynamical model MJO forecasts diverge considerably during Week-2, with some models weakening the signal and others maintaining amplitude over the West Pacific.**
- **Based on recent observations and the dynamical guidance, there is considerable uncertainty regarding the evolution of the MJO during the next two weeks.**
- **The MJO may contribute to enhanced convection over parts of the Maritime Continent during the upcoming week.**

**Additional potential impacts across the global tropics and a discussion for the U.S. are available at:**  
**<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>**




# 850-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly (easterly) anomalies persisted over the equatorial central Indian Ocean (southern India and the eastern Arabian Sea)


Westerly anomalies continued over the western North Pacific, while anomalies remained generally weak near the Date Line.

Westerly anomalies persisted over the eastern Pacific and strengthened over the tropical Atlantic.



# 850-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow  
Easterly anomalies (blue shading) represent anomalous east-to-west flow



Multiple westerly wind bursts were observed across the western Pacific between January and mid-April.

During April, westerly anomalies were generally persistent across the Maritime continent and far western Pacific.

During much of May and June, westerly anomalies were observed over the eastern Pacific and Atlantic.


Westerly anomalies associated with the South Asian monsoon are evident. Other anomalies are generally weak, though westerly, across the Pacific basin.

Longitude



# OLR Anomalies – Past 30 days

**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**  
**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**



During early to mid-June, enhanced (suppressed) convection was observed across the Arabian Sea and along the Meiyu front (South Asia). Large scale convection decreased over the eastern Pacific, although continued tropical cyclone activity was evident.

During mid- to late June, enhanced convection was observed near the Date Line, the Maritime Continent, and much of the North American Monsoon region. Suppressed convection associated with delayed monsoon activity continued over South Asia.


Areas of suppressed (enhanced) convection were observed across the northern Indian Ocean (western Pacific) during late June and early July. Convective anomalies increased over the eastern Pacific, partly associated with renewed tropical cyclone activity.



# Outgoing Longwave Radiation (OLR)

## Anomalies (2.5°N-17.5°N)

Time  
↓



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

Since early January, enhanced convection has propagated slowly eastward from the Maritime Continent to the central Pacific (red box), interrupted periodically by subseasonal variability.

The MJO became more coherent during April, with the subseasonal envelopes of enhanced and suppressed convection modulating the strength of the low frequency signal. The anomalous tropical convection pattern became largely incoherent during mid-May, with convection over the eastern Pacific (red box).

During June, the MJO became more organized as eastward propagation of a coherent large-scale convective pattern was observed. The subseasonal pattern has become less coherent during early July.

Longitude




# 200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time



Longitude

A slow eastward progression of negative anomalies was observed from late December to the present across the Indo-Pacific warm pool region (red box).

During February through April, anomalies propagated eastward with time associated with the MJO before weakening for much of May.

The pattern became more organized during June with a coherent wave-1 structure propagating eastward.


More recently, the pattern became less coherent as other modes of tropical variability (e.g., equatorial Rossby Wave activity) interfered with the subseasonal signal.



# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The upper-level anomalous velocity potential spatial pattern became less coherent during early July, with large scale anomalous ascent partly associated with MJO activity observed over Africa. Rossby Wave activity is evident over the Indian Ocean, and tropical cyclone activity is influencing the pattern over the eastern and western Pacific.




# 200-hPa Vector Wind Anomalies ( $\text{m s}^{-1}$ )

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Upper-level westerly wind anomalies persisted over the equatorial Indian Ocean, with a small area of easterly anomalies developing over the equatorial Atlantic.



# 200-hPa Zonal Wind Anomalies ( $\text{m s}^{-1}$ )

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



Time



Longitude

From December into March, westerly anomalies were most prevalent across the western Hemisphere (red box).


During mid-April, the slowly evolving background state has contributed to easterly anomalies expanding to the Date Line.

MJO activity is evident in the eastward propagation of both easterly and westerly anomalies during April and early May. This signal weakened during late May.

Westward propagation of westerly anomalies is evident over the east central Pacific during June, as is the eastward propagation of a larger scale signal.



# Weekly Heat Content Evolution in the Equatorial Pacific



Oceanic downwelling Kelvin wave activity is evident in late August 2013 and once again during October through early December 2013.

A considerably stronger downwelling event began in January 2014 and propagated across the Pacific.

Warm anomalies persisted over much of the Pacific during April and May, though basin-averaged anomalies weakened during June.



# MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

**Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.**


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

**Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.**

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



# MJO Index -- Recent Evolution




- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The MJO index remained weak during the previous two weeks.



# MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 2007 to present.

Plot puts current MJO activity in recent historical context.




# Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members  
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts  
dark gray shading: 50% of forecasts

The ensemble GFS forecast indicates a strengthening signal with eastward propagation of the MJO RMM Index over the Maritime Continent during the next week. Most ensemble members weaken the signal considerably during Week-2.







# Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The ensemble mean GFS forecasts a development of large scale convective anomalies over the northeastern Indian Ocean, Maritime Continent, and western Pacific during Week-1. Convective anomalies weaken during Week-2.




# Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days





The constructed analog MJO forecast depicts a weak anomaly pattern during the upcoming two weeks.



# MJO Composites – Global Tropics

850-hPa Velocity Potential and  
Wind Anomalies (May-Sep)

Precipitation Anomalies (May-Sep)






# U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



# U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>