



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
October 13, 2014**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- The MJO remained weak during the past week.
- Other types of subseasonal variability, including tropical cyclones, continue to play a large role in determining the pattern of anomalous tropical convection.
- Dynamical and statistical models indicate a strengthening signal during Week-1 with a return to a weak MJO during Week-2. The Week-1 amplification signal is likely related to constructive interference of other modes and tropical cyclone activity.
- Based on recent observations and model guidance, the MJO is not expected to contribute to anomalous convection across the global tropics.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Easterly anomalies diminished over India while some westerly anomalies expanded westward over Southeast Asia.

Easterly anomalies replaced westerly anomalies near the equatorial West Pacific

Anomalies remained weak across the East Pacific.

Westerly anomalies over the Atlantic expanded.




850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time
↓



Longitude

A westerly wind burst was observed across the western Pacific in mid-April.

During April, westerly anomalies were generally persistent across the Maritime continent and far western Pacific.

During much of May and June, westerly anomalies were observed over the eastern Pacific. An enhanced South Asian monsoon circulation developed during much of June and July.

From late July to August, westerly (easterly) anomalies shifted westward over the eastern and central Pacific (western Pacific, Maritime Continent, and Indian Ocean).


Weak easterly (westerly) anomalies developed near the Maritime Continent (Americas), with weak anomalies across much of the Pacific.



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)



During mid-September, enhanced (suppressed) convection was observed over Southeast Asia and Central America (Central Pacific).


During mid to late September, suppressed (enhanced) convection was located over Southeast Asia and East Pacific (West Pacific and parts of North America). Suppressed convection remained over Northern South America.

From late September to early October, suppressed convection became entrenched over Southeast Asia and the Maritime Continent, while tropical cyclone activity contributed to enhanced convection over the West Pacific.



Outgoing Longwave Radiation (OLR)

Anomalies (2.5°N-17.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

Since April, enhanced convection has propagated slowly eastward from the Maritime Continent to the central Pacific (red box).

The anomalous tropical convection pattern became largely incoherent during mid-May. During June and portions of July, the MJO became more organized, primarily over the Indian Ocean, but the pattern became less coherent with respect to canonical MJO activity through August.

There is evidence of westward-moving subseasonal variability from mid-August and later. Some evidence of faster, eastward moving waves is also evident, although nothing consistent with MJO activity on a broad scale.




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time



A slow eastward progression of negative anomalies was observed from March to present across the Indo-Pacific warm pool and central Pacific (red box).

During April, anomalies propagated eastward with time associated with the MJO before weakening for much of May.

The pattern became more organized during June with a more coherent wave-1 MJO-like structure with eastward propagation.

The pattern became less coherent during early July, but then organized again in late July and August, with a wide area of suppressed convection moving around the planet.


During early September, anomalies were consistent with rapid eastward propagation, before becoming stationary for the second half of the month. During October, some eastward propagation is evident over Africa, with a wave-1 structure possibly developing.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The upper-level anomalous velocity potential spatial pattern indicates a wave-1 structure, supportive of enhanced (suppressed) convection from the East Pacific to the Atlantic (Africa to the Maritime Continent).




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Westerly wind anomalies developed along the equator over the eastern Pacific and the Indian Ocean. Easterly anomalies over the Western Pacific also diminished. The Asian Monsoon circulation remained weaker than average as well.



200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow



MJO activity is evident in the eastward propagation of both easterly and westerly anomalies during April and early May. This signal weakened during late May.

Westward propagation of westerly anomalies is evident over the east-central Pacific during June. In July, easterly anomalies intensified over the central and eastern Pacific.


A slow, eastward progression of westerly anomalies is evident over the Maritime Continent and western Pacific during August.

More recently, strong westerly anomalies have developed over the eastern Pacific, with easterly anomalies near 140E.



Weekly Heat Content Evolution in the Equatorial Pacific

Time
↓



Oceanic downwelling Kelvin wave activity is evident during October through early December 2013.

A considerably stronger downwelling event began in January 2014 and propagated across the Pacific.

Warm anomalies persisted over much of the Pacific during April and May, though basin-averaged anomalies decreased during June associated with upwelling Kelvin wave activity (dotted line).

Warm anomalies are again evident across much of the Pacific due to another downwelling Kelvin wave.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution




- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

The RMM MJO index indicates a weak signal with enhanced convection over the Americas.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 2007 to present.

Plot puts current MJO activity in recent historical context.



Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members


Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The ensemble GFS forecast indicates a short lived strengthening of the signal during Week-1, then a nearly stationary signal for Week-2.







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The ensemble mean GFS forecasts a nearly stationary strengthening of the signal during the next 2 weeks.




Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The constructed analog forecast also depicts a weak pattern with some eastward propagation of suppressed convection over the Maritime Continent.




MJO Composites – Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (May-Sep)



Precipitation Anomalies (May-Sep)






U.S. MJO Composites – Temperature

- Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9


<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

▪ Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

▪ Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>