

Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
28 December 2015

Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

Overview

The MJO remained active during the past week, with the enhanced phase now over the West Pacific.

Dynamical model forecasts of the MJO index all support additional eastward propagation across the Pacific during Week-1, with increasing divergence among the forecasts of amplitude and the eastward extent of the signal by late Week-2.

The MJO is constructively interfering with the background El Niño state, resulting in widespread intense convection across the equatorial central Pacific.

The MJO is likely to continue playing a role in the pattern of anomalous convection along with the ongoing El Niño.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>


850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

The coverage of westerly anomalies diminished north of the equator over the Maritime Continent, but persisted south of the equator.



850-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

The red box highlights the persistent low-frequency westerly wind anomalies associated with ENSO.

A robust MJO event was observed in late June through mid-July. Otherwise, tropical cyclone activity across much of the Pacific provided the primary transient influence on the overall ENSO pattern for much of the NH summer.

An eastward shift in the pattern was observed in late October, related to subseasonal activity.

More recently, renewed MJO activity produced an eastward propagation of westerly anomalies from the Indian Ocean to the west-central Pacific during December.



OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During late November and early December, enhanced (suppressed) convection persisted over the Indian Ocean and east-central Pacific (Maritime Continent and tropical South America), consistent with ENSO.

During early to mid-December, the MJO enhanced (suppressed) phase significantly reduced the dry (wet) signal over the Maritime Continent (central Pacific). Suppressed convection persisted over northern South America.

During mid- to late December, enhanced monsoon convection persisted over northern Australia, while suppressed convection returned to the Northwest Pacific. Constructive interference between the MJO and ENSO resulted in widespread enhanced convection across the central Pacific.



Outgoing Longwave Radiation (OLR) Anomalies (5°N-5°S)


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

Since April, the ongoing El Niño is observed (red box) as a tendency toward a dipole of anomalous convection extending from the Maritime Continent (suppressed) to the East Pacific (enhanced).

During June and early July, the MJO become active, interfering with the ENSO signal at times. Since July, the MJO has remained weak, with strong El Niño conditions and tropical cyclone activity dominating the pattern.

The tripole pattern of enhanced/suppressed/enhanced convection stretching from the Indian Ocean to the eastern Pacific has shifted slightly to the east during December.



200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation


Negative anomalies (green shading) indicate favorable conditions for precipitation

The ongoing ENSO state is highlighted by the red box, showing anomalous divergence over the central and eastern Pacific. This pattern has only been temporarily interrupted by strong Kelvin wave/MJO activity at times.


During June and early July, a high-amplitude MJO event was observed, constructively interfering with the El Niño signal in early July.

From July through early October, a generally stationary pattern, reflective of El Niño conditions, was observed. During late October, there was an eastward shift in the pattern associated with subseasonal activity followed by evidence of equatorial Rossby and Kelvin wave activity impacting the central Pacific.

Most recently, strong enhanced divergence expanded east from the Indian Ocean to the central Pacific.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



The upper-level velocity potential anomaly pattern became slightly less coherent, now exhibiting a Wave-2 pattern. A strong couplet of enhanced (suppressed) upper-level divergence is still evident over the Pacific (Indian Ocean and western Maritime Continent).

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation
Negative anomalies (green contours) indicate favorable conditions for precipitation


200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

The ENSO-enhanced subtropical jet weakened west of North America and was suppressed to the south as a highly amplified mid-latitude pattern became established.




200-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Easterly anomalies have persisted over the central and eastern Pacific since June associated with El Niño (red box). During June and July, these easterly anomalies were briefly interrupted by robust atmospheric Kelvin wave/MJO activity.

During late October and again in mid-December, a temporary eastward shift in the westerly anomalies is evident across the Pacific.




Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

Following a strong westerly wind burst in March, a strong downwelling phase of a Kelvin wave propagated eastward, reaching the South American coast during May.

Reinforcing downwelling events have followed, resulting in persistently above-normal heat content from the Date Line to 80W throughout the period.

An expansion of below average heat content over the western Pacific is evident since spring and this area has increased during November and December 2015.



MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


The MJO index remained highly amplified during the past two weeks, with fast propagation from the Maritime Continent to the West Pacific.



MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



Ensemble GFS (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The GFS ensemble MJO index forecast depicts a continuation of the eastward propagation over the West and Central Pacific during the next two weeks.


Yellow Lines - 20 Individual Members
Green Line - Ensemble Mean



Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast
Initial date: 27 Dec 2015
OLR




The GEFS MJO index-based OLR forecast depicts continued eastward propagation of the intraseasonal convective anomalies, with increased magnitude during Week-2.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:27-Jun-2015 to 27-Dec-2015
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model
reconstruction by RMM1 & RMM2 (27 Dec 2015)




The constructed analog model also depicts eastward propagation of strong MJO-related convective anomalies.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S- 7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5° S, 7.5° N] (cont:4Wm⁻²) Period:27-Jun-2015 to 27-Dec-2015
The unfilled contours are CA forecast reconstructed anomaly for 15 days




MJO Composites - Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov-Mar)




Precipitation Anomalies (Nov-Mar)



U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>