

Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
14 March 2016

Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

Overview

The enhanced phase of the MJO entered the Indian Ocean during the past week with its suppressed phase destructively interfering with the ongoing El Nino across the Pacific.

The dynamical model MJO index forecasts support a continued eastward propagation although differences in strength of the MJO signal exist.

Destructive interference between the intraseasonal signal and the ENSO background state is likely during the next week.

During Week-1, the MJO favors enhanced convection across northern Australia with increased chances of tropical cyclone formation over the southern Indian Ocean and Coral Sea region. Anomalous convection associated with ENSO is expected to persist but potentially be offset by the MJO signal.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>


850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Easterly anomalies diminished across the Indian Ocean during the past five days with westerly anomalies expanding across the southern Indian Ocean.



Easterly (westerly) anomalies persisted across the eastern Maritime Continent (Western Hemisphere).

850-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

The red box highlights the persistent low-frequency westerly wind anomalies associated with ENSO.

An eastward shift in the pattern was observed in late October, related to subseasonal activity.

MJO activity during December produced an eastward propagation of westerly anomalies from the Indian Ocean, which constructively interfered with El Niño during January, and lead to a westerly wind burst near the Date Line. Another period of constructive interference occurred in late February.

Most recently, easterly anomalies have overspread the Maritime Continent and westerly anomalies are increasing over the eastern Pacific.



OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

The pattern of anomalous convection was generally consistent with ENSO during mid-February although suppressed convection was disrupted across the western Maritime Continent.

During late February, the MJO constructively interfered resulting in suppressed (enhanced) convection across the Maritime Continent and northern Australia (equatorial central Pacific).

Convection increased across northern South America during early March as the enhanced phase of the MJO propagated across the Western Hemisphere.



Outgoing Longwave Radiation (OLR) Anomalies (17.5°S-2.5°S)

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)


Wetter-than-normal conditions, negative OLR anomalies (blue shading)

Since April, the ongoing El Niño is observed (red box) as a dipole of anomalous convection extending from the Maritime Continent to the East Pacific.

During December, the MJO became active again, with the enhanced phase propagating from the Indian Ocean to the west-central Pacific during the month.

Renewed MJO activity was evident during late January and February, with enhanced convection propagating over the Indian Ocean and a reduction of suppressed convection over the Maritime Continent.

Recently, the MJO is contributing to an eastward shift on convection over the eastern Pacific and suppressed rainfall over the western Pacific.



200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation


The ongoing ENSO state is highlighted by the red box, showing anomalous divergence over the central and eastern Pacific.

During June and early July, a high-amplitude MJO event was observed, constructively interfering with the El Niño signal in early July.


During late October, there was an eastward shift in the pattern. Renewed MJO activity was also observed during December and early January, yielding a robust signal in the upper levels. This signal weakened during mid-January.

During late February, intraseasonal variability constructively interfered with the ongoing El Niño.

Recently, upper-level divergence entered the Indian Ocean.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



The velocity potential anomalies depict upper-level divergence entering the Indian Ocean with the area of upper-level convergence disrupted by the background ENSP state over the Pacific Ocean.

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation
Negative anomalies (green contours) indicate favorable conditions for precipitation

200-hPa Vector Wind Anomalies (m s^{-1})


Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Easterly (westerly) anomalies persisted over the eastern and central Pacific (Indian Ocean).

A large subtropical High (common during El Niño events) has redeveloped over the eastern Pacific.



200-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Easterly anomalies have persisted over the central and eastern Pacific since June associated with El Niño (red box).

During late October, a temporary eastward shift in the westerly anomalies was evident across the Pacific.

Eastward propagation of upper-level zonal wind anomalies was apparent over the Maritime Continent and West Pacific during late December and early January, consistent with MJO activity.

More recently, westerly anomalies returned to the Indian Ocean and Maritime Continent, while easterly anomalies persisted between about 170E - 120W.




Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

Following a strong westerly wind burst in March, a strong downwelling phase of a Kelvin wave propagated eastward, reaching the South American coast during May.

Reinforcing downwelling events have followed, resulting in persistently above-normal heat content from the Date Line to 80W throughout the period.

An eastward expansion of below average heat content over the western Pacific is evident since January, and negative anomalies spread east of the Date Line during February 2016. The below average heat content has also increased in magnitude.



MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


The MJO index indicates continued eastward propagation to the Indian Ocean during the early March.



MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



Ensemble GFS (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The GFS ensemble MJO index forecast depicts a continuation of MJO activity during mid-March with apparent destructive interference with the background ENSO state.


Yellow Lines - 20 Individual Members
Green Line - Ensemble Mean



Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast
Initial date: 13 Mar 2016
OLR




The GEFS OLR forecast depicts enhanced convection shifting east across the Maritime Continent during the next two weeks.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S– 7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5° S, 7.5° N] (cont: 4Wm^{-2}) Period: 12-Sep-2015 to 13-Mar-2016
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model
reconstruction by RMM1 & RMM2 (13 Mar 2016)




The constructed analog model also depicts eastward propagation of the MJO-associated OLR anomalies.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5°S,7.5°N] (cont:4Wm⁻²) Period:12-Sep-2015 to 13-Mar-2016
The unfilled contours are CA forecast reconstructed anomaly for 15 days




MJO Composites - Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (Nov-Mar)




Precipitation Anomalies (Nov-Mar)



U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>