

# Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:  
Climate Prediction Center / NCEP  
27 February 2017

# Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

# Overview

- The MJO is active, with both the RMM based and CPC velocity potential based MJO indicating enhanced (suppressed) convection over the Atlantic, Africa, and the Indian Ocean (Pacific Ocean).
- The MJO is constructively interfering with the low frequency state over the Pacific basin.
- Dynamical model RMM index forecasts indicate a reduction in amplitude and some eastward propagation in Week-1, followed by a weak signal in Week-2. Statistical model outputs are similar to the dynamical model outputs, but maintain higher amplitudes during Week-2.
- The MJO is anticipated to influence the evolution of the global tropical convective pattern over the next several days. During Week-1, some constructive interference between the MJO and the base state is likely over the Pacific, while uncertainty increases in Week-2. Impacts during the next two weeks to the extratropics remain uncertain given the interaction between the MJO and an evolving low frequency state.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:  
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>


# 850-hPa Vector Wind Anomalies (m s<sup>-1</sup>)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Easterly anomalies persisted over the southern Indian Ocean and expanded across the Maritime Continent and West Pacific, consistent with ongoing MJO activity.



Westerly anomalies remained prominent over the East Pacific and the Caribbean Sea.

# 850-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)


Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Persistent westerly (easterly) anomalies were evident over the eastern Indian Ocean and western Maritime Continent (central and western Pacific) as shown by the red (blue) box at right. These anomalies are low frequency in nature, associated with the negative phase of the Indian Ocean Dipole (IOD), and later, La Niña.

During mid-January, Rossby Wave activity was evident, with destructive interference on the base state evident through 100E.

Over the past couple of weeks, eastward propagating anomalies were observed, consistent with ongoing MJO activity.



# OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

From late January to early February, convection was enhanced (suppressed) over the Maritime Continent and western Pacific (Central Pacific and Indian Ocean).

During early to mid-February, the intraseasonal signal began to interfere with the low frequency state. Enhanced convection developed over the South Pacific, and decreased in magnitude over the Maritime Continent.

During mid- to late February, the MJO continued eastward propagation with enhanced convection developing over parts of Africa and the western Indian Ocean. Suppressed convection was observed over much of the southern Maritime Continent.



# Outgoing Longwave Radiation (OLR) Anomalies (2.5°S - 17.5° S)


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

A low frequency state favoring enhanced convection over the eastern Indian Ocean and the Maritime Continent has been evident since July (green box), with suppressed convection over the Indian Ocean and near the antimeridian (black boxes).

An intraseasonal event occurred during late November and early December that interfered with the background state.

Since late January, an active MJO has shifted the low frequency pattern eastward, with enhanced convection shifting across the Pacific and suppressed convection moving toward the Maritime Continent. More recently, the MJO has led to enhanced convection over parts of the Indian Ocean.



# 200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation


In early September, intraseasonal activity was apparent, before reversion to the low frequency pattern associated with the negative IOD and La Niña through late October.

During November, eastward propagation was observed consistent with MJO activity on the fast end of the intraseasonal spectrum.


After a break in apparent MJO activity during December, a signal emerged over the Maritime Continent and has continued propagating through the present.

There have been alternating periods of constructive and destructive interference between the MJO and the low-frequency state. Most recently there has been constructive interference over the Pacific.

200-hPa Velocity Potential Anomaly: 5N-5S



# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



Upper-level velocity potential anomalies reveal a wavenumber-1 pattern. Enhanced divergence is evident from the Atlantic to the Indian Ocean and western Maritime Continent, with enhanced subsidence centered over the Pacific basin.

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation  
Negative anomalies (green contours) indicate favorable conditions for precipitation

# 200-hPa Vector Wind Anomalies (m s<sup>-1</sup>)


Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Anomalous easterlies have been fairly persistent over the East Pacific.

There appears to have been a rapid reversal of anomalies related to the extension/retraction of the East Asian jet stream from the early period to the most recent pentad.



# 200-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Eastward propagation of westerly anomalies was broadly consistent with organized MJO activity during September.

In November, anomalous westerlies persisted near the Date Line, though intraseasonal variability associated with the MJO is evident.

In late November, easterly anomalies re-emerged across the Indian Ocean and Maritime Continent, consistent with the passage of sub-seasonal activity and the re-alignment of the low frequency base state.

Near the end of 2016 a period of westerlies disrupted the low frequency state between 80-130E and continued propagating eastward through the Western Hemisphere. This subseasonal activity has continued, with alternating anomalous westerlies/easterlies being observed over the Pacific.




# Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

An eastward expansion of below average heat content over the western Pacific is evident through June, with widespread negative anomalies building across the Pacific over the course of boreal spring and summer.

More recently, upper-ocean heat content anomalies have been low amplitude, consistent with the forecast transition to ENSO-neutral conditions. Positive anomalies are now observed over the entire basin.



# MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

# MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes

The triangular areas indicate the location of the enhanced phase of the MJO


Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months

During the past week, continued eastward propagation of an MJO signal was evident across Africa to the western Indian Ocean.


[RMM1, RMM2] Phase Space for 18-Jan-2017 to 26-Feb-2017



# MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



# GFS Ensemble (GEFS) MJO Forecast

RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days


light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The GEFS depicts some eastward propagation along with dramatic reduction in amplitude during Week-1.


The displacement towards Phases 7 and 8 of a counterclockwise propagating signal in RMM phase space could be a result of a changing base state with conditions becoming more favorable for convection in the Central Pacific.

Yellow Lines - 20 Individual Members  
Green Line - Ensemble Mean



# Ensemble GFS (GEFS) MJO Forecast


Spatial map of OLR anomalies for the next 15 days



The GEFS prediction for RMM Index-based OLR anomalies over the next two weeks shows some eastward propagation and gradual weakening of enhanced convection across the Indian Ocean.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days



# Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (26 Feb 2017)




The statistical (Constructed Analog) RMM-based OLR anomaly prediction shows similar propagation to the dynamical model guidance, but at a much higher amplitude late in the period.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cont:4Wm<sup>-2</sup>) Period:27-Aug-2016 to 26-Feb-2017  
The unfilled contours are CA forecast reconstructed anomaly for 15 days




# MJO Composites - Global Tropics

850-hPa Velocity Potential and  
Wind Anomalies (Nov - Mar)




Precipitation Anomalies (Nov - Mar)



# U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

# U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>