

# Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:  
Climate Prediction Center / NCEP  
20 March 2017

# Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

# Overview

- The MJO exhibited continued weakness during the past week, as a unique scenario with two low-frequency centers of action has emerged over the East Pacific and Maritime Continent.
- Dynamical model RMM index forecasts generally depict a weak MJO signal during the next two weeks generally centered towards the Indian Ocean. This solution is discounted, due to apparent influences of the multiple areas of active convection and inconsistent placement of any potential subseasonal activity.
- Given the unique, robust low frequency footprint, the MJO is not anticipated to play a major role in the evolution of the global tropical convective pattern during the next two weeks.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:  
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>


# 850-hPa Vector Wind Anomalies (m s<sup>-1</sup>)

Note that shading denotes the zonal wind anomaly

**Blue shades:** Easterly anomalies

**Red shades:** Westerly anomalies

Weak, spatially variable low-level anomalies were spread over the Indian Ocean and Maritime Continent.



Easterly anomalies persisted over the eastern Maritime Continent and West Pacific.

Westerly anomalies persisted over the far East Pacific and equatorial South America.

# 850-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Persistent westerly (easterly) anomalies were evident over the eastern Indian Ocean and western Maritime Continent (central and western Pacific) as shown by the red (blue) box at right. These anomalies are low frequency in nature, associated with the negative phase of the Indian Ocean Dipole (IOD), and later, La Niña.

During late January, Rossby Wave activity was evident, with destructive interference on the base state evident through 100E.

During February, eastward propagating anomalies were observed, consistent with ongoing MJO activity.

More recently, other modes interfered with the intraseasonal signal. Westward moving features are evident from the date line through the Maritime Continent.



# OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During mid and late February, enhanced (suppressed) convection was observed over Africa, the northwest Pacific, South America and tropical Atlantic (southern Indian Ocean, Central and eastern North Pacific)

As the intraseasonal signal returned to the Indian Ocean in late February and early march, enhanced convection overspread the northern and equatorial Indian Ocean and interacted with a westward moving feature over the Maritime Continent.

During early to mid-March, tropical cyclone activity was evident over the southern Indian Ocean. Influence from Rossby wave activity and the remnant low frequency signal became increasingly apparent, resulting in a weakening intraseasonal signal.



# Outgoing Longwave Radiation (OLR) Anomalies (2.5°S - 17.5° S)

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)


Wetter-than-normal conditions, negative OLR anomalies (blue shading)

A low frequency state favoring enhanced convection over the eastern Indian Ocean and the Maritime Continent has been evident from July through Mid-February (green box), with suppressed convection over the Indian Ocean and near the antimeridian (black boxes).

An intraseasonal event during late November and early December interfered with the background state.

Since late January, an active MJO pattern became the dominant mode of intraseasonal tropical convective variability, with the suppressed phase reversing the low frequency enhanced convective signal over the Maritime Continent in late February.

The MJO envelope may be present near the Maritime Continent recently, although this could be the lingering low frequency state.



# 200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation


In late September, intraseasonal activity was apparent, before reversion to the low frequency pattern associated with the negative IOD and La Niña through late October.

During November, eastward propagation was observed consistent with MJO activity on the fast end of the intraseasonal spectrum.


After a break in apparent MJO activity during December and early January, a signal emerged over the Maritime Continent and has continued propagating through early March.

There have been alternating periods of constructive and destructive interference between the MJO and the low frequency state. Most recently, the remnant intraseasonal signal has begun destructively interfering with the base state over the central Pacific.

200-hPa Velocity Potential Anomaly: 5N-5S  
5-day Running Mean



# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



A wave-2 pattern is evident, with enhanced convective centers of action over the East Pacific and Maritime Continent (suppressed conditions across the Central Pacific, Africa, and the Atlantic). This pattern appears most consistent with the low-frequency background state, with some intraseasonal influence possible across Africa and the South Pacific.

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation

# 200-hPa Vector Wind Anomalies (m s<sup>-1</sup>)


Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

The Northern Hemisphere extratropics remained fairly consistent over the past two pentads, with the exception of recent wave breaking across Central U.S. and Atlantic.

Anomalous easterlies persisted just south of the equator in the East Pacific, associated with the anomalous convection in the Niño 1+2 region.



# 200-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Eastward propagation of westerly anomalies was broadly consistent with organized MJO activity during September.

In November, anomalous westerlies persisted near the Date Line, though intraseasonal variability associated with the MJO is evident.

In late November, easterly anomalies re-emerged across the Indian Ocean and Maritime Continent, consistent with the passage of sub-seasonal activity and the realignment of the low frequency base state.

Near the end of 2016 a period of westerlies disrupted the low frequency state between 80-130E and continued propagating eastward through the Western Hemisphere. This subseasonal activity has continued, with alternating anomalous westerlies/easterlies being observed over the Pacific.




# Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

An eastward expansion of below average heat content over the western Pacific is evident through June, with widespread negative anomalies building across the Pacific over the course of boreal spring and summer.

More recently, upper-ocean heat content anomalies have been low amplitude, consistent with the forecast transition to ENSO-neutral conditions. Positive anomalies are now observed over the entire basin.



# MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

# MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


During the past week, the RMM index continued to indicate weak MJO activity.



# MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



# GFS Ensemble (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The GEFS depicts a weak MJO signal during Week-1, with the potential emergence of a weak, stagnant signal over the Indian Ocean during Week-2.


Yellow Lines - 20 Individual Members  
Green Line - Ensemble Mean



# Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast  
Initial date: 19 Mar 2017  
OLR




The GEFS prediction for RMM Index-based OLR anomalies over the next two weeks shows a weak anomaly pattern early in the period with stationary amplification throughout the next two weeks.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2  
OLR [7.5°S,7.5°N] (cont:4Wm<sup>-2</sup>) Period:17-Sep-2016 to 19-Mar-2017  
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



# Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (19 Mar 2017)




The statistical (Constructed Analog) RMM-based OLR anomaly prediction shows minimal eastward propagation of an MJO signal with limited change in intensity.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cont:4Wm<sup>-2</sup>) Period:17-Sep-2016 to 19-Mar-2017  
The unfilled contours are CA forecast reconstructed anomaly for 15 days




# MJO Composites - Global Tropics

850-hPa Velocity Potential and  
Wind Anomalies (Nov - Mar)




Precipitation Anomalies (Nov - Mar)



# U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

# U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>