

Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
26 June 2017

Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

Overview

- The MJO remains weak, and is not a major contributor to convection across the tropics.
- Dynamical model guidance varies on the evolution of the MJO during the next two weeks. Models generally increase the convection over the Indian Ocean during Week-1, but then some models move that signal westward, while others move it eastward, so uncertainty in Week-2 is high.
- The MJO is expected to strengthen during Week-1, then weaken slightly during week-2, with the primary impacts being over the Indian Ocean basin.


Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>

850-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Monsoon flow over India remained weaker than average, though closer to normal.

Easterly (Westerly) wind anomalies persisted over the western (eastern) Pacific.

850-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Persistent westerly (easterly) anomalies, shown by the red (blue) box at right, were associated with the negative phase of the Indian Ocean Dipole (IOD), and later, La Niña.

During late January, Rossby wave activity was evident, with destructive interference on the base state evident through 100E.

During February, MJO activity also destructively interfered with the base state. During mid-March and early April, the low frequency state reemerged, with some intraseasonal variability evident in late March.

Most recently, westerly (easterly) anomalies became more coherent across the Western (Eastern) Hemisphere. Recently, equatorial anomalies have been weak.



OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

During late May into the beginning of June, enhanced convection was apparent over the eastern Maritime Continent, with suppressed conditions persisting over much of the remainder of the global tropics.

A robust Kelvin wave increased convection across parts of the East Pacific, northern South America, the tropical Atlantic, and Africa during early June.

During mid-June, the pattern became stagnant, with enhanced precipitation remaining over Africa and portions of the Caribbean. The South Asian Monsoon waned and dry conditions broadly remained over portions of the western Pacific.



Outgoing Longwave Radiation (OLR) Anomalies (2.5°N - 17.5°N)


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

A low frequency state favoring enhanced convection over the eastern IO and the Maritime Continent was evident from July 2016 through early April 2017 (green box), with suppressed convection near the Date Line (right black box). The remainder of the IO generally had suppressed convection during this period (left black box), with the exception of an MJO-related wet period from mid-Feb to early March.

From mid-April through present, convective anomalies were generally weak. In mid-May, enhanced convection was noted over the Indian Ocean with some eastward propagation.

The most recent pronounced signal is related to convective suppression between 80-160W.



200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation


The pattern, during December and January, was more related to seasonal variability.

A signal emerged over the Maritime Continent and continued propagating through early March, creating alternating periods of constructive and destructive interference with the base state.


During March, a low frequency signal favoring enhanced (suppressed) convection over the Maritime Continent (Indian Ocean) once again became the primary component of the anomaly field.

Kelvin wave activity has been apparent from April through the present, evident in the rapidly propagating eastward signals. During later June, anomalies in velocity potential have become almost stationary.

200-hPa Velocity Potential Anomaly: 5N-5S
5-day Running Mean



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



Multiple waves are evident and the pattern is not consistent with robust MJO activity.

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation


200-hPa Vector Wind Anomalies (m s⁻¹)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Anomalies along the equator are relatively weak, having weakened since last week.



200-hPa Zonal Wind Anomalies (m s⁻¹)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Near the end of 2016 a period of westerlies disrupted the low frequency state between 80-130E and continued propagating eastward through the Western Hemisphere.

Easterly anomalies returned to the East Pacific during late April.

During early to mid-June, easterly anomalies were most prominent across the global tropics, in part due to mid-latitude influences.

Recently, anomalies along the equator are relatively weak, with fast moving eastward propagating signals .



MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months


During the past week, the RMM index decreased in amplitude, reflecting almost no signal during the past 5 days.



MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



GFS Ensemble (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The GEFS forecast predicts no coherent MJO signal during the next 2 weeks, with convection remaining over Africa and the western Indian Ocean.


Yellow Lines - 20 Individual Members
Green Line - Ensemble Mean



Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast
Initial date: 25 Jun 2017
OLR




The GEFS RMM-based OLR anomaly forecasts enhanced (suppressed) convection persisting over Africa and much of the Indian Ocean (West Pacific). By Week-2, the signal shows some westward propagation to enhance convection over the Americas, as well.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2
OLR [7.5°S,7.5°N] (cint:4Wm⁻²) Period:24-Dec-2016 to 25-Jun-2017
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (25 Jun 2017)




The constructed analog RMM-based OLR anomaly prediction indicates little to no propagation of a weak signal, with convection remaining over the Indian Ocean.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [7.5°S,7.5°N] (cint:4Wm²) Period:24-Dec-2016 to 25-Jun-2017
The unfilled contours are CA forecast reconstructed anomaly for 15 days



U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>