

# Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:  
Climate Prediction Center / NCEP  
7 August 2017

# Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

# Overview

- The MJO weakened rapidly over the past several days as the large-scale convective envelope over the Maritime Continent and West Pacific collapsed due in part to interference from a highly amplified mid-latitude pattern over the Southern Hemisphere.
- A weak remnant signal is evident over the central Pacific, and a Kelvin wave is currently propagating across the Indian Ocean.
- Dynamical and statistical model guidance depict a weak MJO signal over the next two weeks, with potential tropical cyclone activity over the East Pacific also influencing the signal. Longer range CFS forecasts suggest potential renewed MJO activity over the Indian Ocean, during either late Week-2 or into the Week-3 period.
- The MJO is not anticipated to influence the global tropical convective pattern during the next two weeks.


Additional potential impacts across the global tropics and a discussion for the U.S. are available at:  
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>

# 850-hPa Vector Wind Anomalies (m s<sup>-1</sup>)

Note that shading denotes the zonal wind anomaly

**Blue shades:** Easterly anomalies

**Red shades:** Westerly anomalies



A weakened monsoon signal is very apparent in the low-latitude zonal wind anomalies over South Asia.

An amplified mid-latitude pattern is evident over the southern Indian Ocean and Maritime Continent, with some potential influence on the tropics.

Weak easterlies (westerlies) persist over the central Pacific (Atlantic and equatorial Indian Ocean), reflecting the remnant interaction of the intraseasonal and low-frequency signals.

# 850-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Persistent westerly (easterly) anomalies, shown by the red (blue) box at right, were associated with the negative phase of the Indian Ocean Dipole (IOD), and La Niña.

In February, MJO activity destructively interfered with the base state. During mid-March and early April, the low frequency state reemerged, with some intraseasonal variability evident in late March.

Equatorial flow was fairly weak throughout June, with easterlies favored between 120E and the Date Line and also emerging across the western Indian Ocean.

Since late June, the low-frequency pattern shifted eastward, with easterly anomalies over the central Pacific. More recently, westerly anomalies strengthened over the Indian Ocean while the overall pattern weakened.



# OLR Anomalies - Past 30 days


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

In early July, large-scale enhanced convection was observed over the Maritime Continent, surrounded by suppressed convection (western Indian Ocean and central Pacific). TCs and enhanced monsoon flow were evident over the East Pacific and southwest US.

During mid-July, slow eastward propagation of the large-scale anomalies was evident.

By late July and early August, the enhanced convective signal over the Maritime Continent broke down. Strong TC activity over the northwestern Pacific is evident. The South Asian Monsoon pattern weakened considerably, with suppressed (enhanced) convection over India (central Indian Ocean just south of the Equator).



# Outgoing Longwave Radiation (OLR) Anomalies (2.5°N - 17.5°N)


**Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)**

**Wetter-than-normal conditions, negative OLR anomalies (blue shading)**

A low frequency state favoring enhanced convection over the eastern IO and the Maritime Continent was evident from July 2016 through early April 2017 (green box), with suppressed convection near the Date Line (right black box).

From mid-April through present, convective anomalies were generally weak. In mid-May, enhanced convection was noted over the Indian Ocean with some eastward propagation.

During mid-July, there was a burst of enhanced convection over the Maritime Continent, due to interactions between a potential intraseasonal signal and the low-frequency state. More recently, the convection waned considerably, with suppressed convection noted over much of South and Southeast Asia.



# 200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation


A signal emerged over the Maritime Continent and continued propagating through early March, creating alternating periods of constructive and destructive interference with the base state.

During March, a low frequency signal favoring enhanced (suppressed) convection over the Maritime Continent (Indian Ocean) once again became the primary component of the anomaly field.


Kelvin wave activity was apparent from April through early June, as seen in the rapidly propagating eastward signals.

During July, enhanced convection strengthened over the Maritime Continent as the low-frequency signal constructively interfered with an easterly propagating signal. More recently, suppressed convection overspread the Maritime Continent, with weakly enhanced convection (Kelvin wave-related) now over the central Pacific.

200-hPa Velocity Potential Anomaly: 5N-5S  
5-day Running Mean



# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



A Wave-2 pattern is evident, with an enhanced (suppressed) signal over the central Pacific (Maritime Continent) and Indian Ocean, and a second, weaker couplet over the Africa and the Americas.

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation


Negative anomalies (green contours) indicate favorable conditions for precipitation

# 200-hPa Vector Wind Anomalies (m s<sup>-1</sup>)

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies




Upper-level convergence is evident over the Maritime Continent. Weak westerly anomalies are present from the Date Line to the coast of Ecuador.



# Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.



Upper-ocean heat content anomalies remain weak across the Pacific, with a sign change (positive to negative) observed over the central Pacific in late July.

# MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

# MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes


The triangular areas indicate the location of the enhanced phase of the MJO

Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months

[RMM1, RMM2] Phase Space for 28-Jun-2017 to 06-Aug-2017




The MJO signal weakened considerably during the past few days as other modes interfered with the intraseasonal signal.

# MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



# GFS Ensemble (GEFS) MJO Forecast


RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days

light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

The GEFS ensembles generally depict a weak MJO signal over the next two weeks.


Yellow Lines - 20 Individual Members  
Green Line - Ensemble Mean



# Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast  
Initial date: 06 Aug 2017  
OLR




The GEFS RMM-based OLR anomaly forecast shows little signal over the next two weeks.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days


Reconstructed anomaly field associated with the MJO using RMM1 & RMM2  
OLR [7.5°S,7.5°N] (cont:4Wm<sup>-2</sup>) Period:04-Feb-2017 to 06-Aug-2017  
The unfilled contours are GEFS forecast reconstructed anomaly for 15 days



# Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (06 Aug 2017)




The constructed analog RMM-based forecast also depicts a weak signal, and suggests a continuation of the suppressed South Asian monsoon signal during Week-2.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of ( $7.5^{\circ}$  S- $7.5^{\circ}$  N) OLR anomalies - last 180 days and for the next 15 days

Reconstructed anomaly field associated with the MJO using RMM1 & RMM2 OLR [ $7.5^{\circ}$ S, $7.5^{\circ}$ N] (cont:4Wm<sup>-2</sup>) Period:04-Feb-2017 to 06-Aug-2017  
The unfilled contours are CA forecast reconstructed anomaly for 15 days






# U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

# U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>