

# Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions



Update prepared by:  
Climate Prediction Center / NCEP  
16 October 2017

# Outline

Overview

Recent Evolution and Current Conditions

MJO Index Information

MJO Index Forecasts

MJO Composites

# Overview

- There has been a recent strengthening of the MJO signal during the past 2 weeks, with the enhanced convective phase located over the Maritime Continent region.
- Available 200-hPa velocity potential anomaly forecasts predict eastward propagation of the current MJO signal from the Maritime Continent to the central Pacific during the next two weeks; thereafter, followed by rapid weakening of the signal.
- The MJO signal favors tropical cyclone development over the West Pacific during Week-1.

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:  
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>


# 850-hPa Vector Wind Anomalies (m s<sup>-1</sup>)

Note that shading denotes the zonal wind anomaly

**Blue shades:** Easterly anomalies

**Red shades:** Westerly anomalies

Westerly anomalies expanded over the tropical Indian Ocean and Maritime Continent, pushing the leading edge of easterly anomalies to about 150E.



Easterly anomalies strengthened just to the west of the Date Line.

Westerly anomalies in the East Pacific weakened substantially during the past week.

# 850-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

Low-frequency easterly anomalies (blue box) have largely persisted over the west-central Pacific throughout the last 180 days.

Equatorial zonal wind anomalies were of low amplitude in June. During July, a slight eastward shift in the low-frequency pattern is noted, related to short-lived MJO activity.

During September, easterly anomalies persisted along and to the west of the Date Line. Some intraseasonal variability is evident, with little to no MJO contribution.

Anomalous westerlies present in the East Pacific during the past few weeks (associated with an enhanced monsoon circulation over Central America), reversed sign about one week ago. An emerging MJO signal is evident with westerlies expanding over the Indian Ocean.



# OLR Anomalies - Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)


Wetter-than-normal conditions, negative OLR anomalies (blue shading)

The persistent area of suppressed convection near the Date Line across the equatorial Pacific is associated with the low frequency state.

Anomalous convection has varied across the Indian Ocean during the past month, while enhanced convection returned to the Maritime Continent. Hurricane Maria's track is apparent in the Atlantic.

More recently, anomalous convection is noted over the Caribbean and western Atlantic, with some contribution from Hurricane Nate. Convective signals weakened over the Maritime Continent once more.

Enhanced convection is noted over parts of the equatorial and North Indian Ocean, and the far western Atlantic; with a disrupted OLR pattern apparent over the Maritime Continent.



# Outgoing Longwave Radiation (OLR) Anomalies (7.5°S - 7.5°N)


Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

In mid-May, enhanced convection was noted over the Indian Ocean with some eastward propagation.

During mid-July, there was a burst of enhanced convection over the Maritime Continent, due to interactions between a short-lived intraseasonal signal and the low-frequency state.

Multiple modes of variability including tropical cyclones contributed to the pattern of anomalous convection during the past month. Suppressed convection continues near the Date Line, while enhanced convection over the Maritime continent remains variable.



# 200-hPa Velocity Potential Anomalies (5°S - 5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation


Kelvin wave activity was apparent from April through early June, as seen in the rapidly propagating eastward signals.

During July, enhanced convection strengthened over the Maritime Continent as the low-frequency signal constructively interfered with an easterly propagating signal. This eastward propagating signal appears more or less intact with a period in line with canonical MJO phase speeds.


A signal on the MJO timescale is evident in this field during late August and September. Some evidence of the suppressed envelope associated with this event is currently analyzed near the Date Line.

Most recently, there is constructive interference between an MJO envelope and the low-frequency state over the Maritime Continent region.

200-hPa Velocity Potential Anomaly: 5N-5S  
5-day Running Mean



# IR Temperatures (K) / 200-hPa Velocity Potential Anomalies



The 200-hPa velocity potential field is wave-1 with enhanced (suppressed) convection over the Eastern (Western) Hemisphere, consistent with MJO presence.

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation

# 200-hPa Vector Wind Anomalies (m s<sup>-1</sup>)


Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies

Anomalous divergence developed over the northern Indian Ocean in early October.

Anomalous westerlies were noted over much of the West Pacific during the last two weeks, increasing in amplitude.



# 200-hPa Zonal Wind Anomalies (m s<sup>-1</sup>)

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow


Easterly anomalies (blue shading) represent anomalous east-to-west flow

During early to mid-June, easterly anomalies were most prominent across the global tropics, in part due to mid-latitude influences.

Starting in July, the anomaly patterns have been continually moving eastward associated with weak MJO activity and atmospheric Kelvin waves.

During September, fast-moving eastward propagation of anomalies continued, consistent with additional atmospheric Kelvin Waves. A slower signal was evident over the eastern Maritime Continent and west Pacific before decaying near 160 W.


Anomalous westerlies gave way to anomalous easterlies during the past week over the Maritime Continent region.



# Weekly Heat Content Evolution in the Equatorial Pacific

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

Upper-ocean heat content anomalies remained about the same amplitude as last week, with some areas warming from 2 weeks ago, highlighting the intraseasonal fluctuations. Overall, OHC is still reflective of the emerging background state.



# MJO Index -- Information

The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.

The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).

# MJO Index - Recent Evolution

The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes

The triangular areas indicate the location of the enhanced phase of the MJO


Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.

Distance from the origin is proportional to MJO strength

Line colors distinguish different months

## PLEASE NOTE:


Due to an upstream data issue, the forecast plot displayed here unfortunately will not be available until close-of-business Oct 17<sup>th</sup> at the earliest. The problem is being worked on, and we very much regret the inconvenience this causes.



# MJO Index - Historical Daily Time Series

Time series of daily MJO index amplitude for the last few years.

Plot puts current MJO activity in recent historical context.



# GFS Ensemble (GEFS) MJO Forecast

RMM1 and RMM2 values for the most recent 40 days and forecasts from the GFS ensemble system (GEFS) for the next 15 days


light gray shading: 90% of forecasts

dark gray shading: 50% of forecasts

## PLEASE NOTE:

Due to an upstream data issue, the forecast plot displayed here unfortunately will not be available until close-of-business Oct 17<sup>th</sup> at the earliest. The problem is being worked on, and we very much regret the inconvenience this causes.


Yellow Lines - 20 Individual Members  
Green Line - Ensemble Mean



# Ensemble GFS (GEFS) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

Prediction of MJO-related anomalies using GEFS operational forecast  
Initial date: 14 Oct 2017  
OLR



Not available today.

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)


Time-longitude section of (7.5° S-7.5° N) OLR anomalies - last 180 days and for the next 15 days

**The GEFS plot of MJO related OLR anomalies is unavailable at this time.**

# Constructed Analog (CA) MJO Forecast

Spatial map of OLR anomalies for the next 15 days

OLR prediction of MJO-related anomalies using CA model  
reconstruction by RMM1 & RMM2 (14 Oct 2017)



Not available today.


Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Time-longitude section of (7.5° S–7.5° N) OLR anomalies - last 180 days and for the next 15 days


**The GEFS plot of MJO related OLR anomalies is unavailable at this time.**

# MJO Composites - Global Tropics

850-hPa Velocity Potential and  
Wind Anomalies (Nov - Mar)




Precipitation Anomalies (Nov - Mar)



# U.S. MJO Composites - Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>

# U.S. MJO Composites - Precipitation

Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>